



James S. Kercher, M.D.
Sports Medicine, Shoulder &
Knee Surgery, Cartilage Restoration

Stable Proximal Humerus Fracture

- This protocol is for nondisplaced proximal humerus fractures, or patients stable fractures after surgical fixation.

Goal: Maintain range of motion (ROM) without displacing the proximal humerus fracture.

Phase I (0 – 21 days)

- Begin elbow, wrist and hand active ROM.
- A sling and swath, or Velpeau with and axillary pad should be used when not doing physical therapy.
- Begin pendulum exercises (clockwise and counterclockwise).
- After 7 days, begin supine ER with a cane. Fifteen to 20 degrees of abduction are permitted if the patient is more comfortable.
- Establish a home exercise program so patient is performing exercises 3-5 times per day for 30 min. each session.

Phase II (3 - 6 weeks)

- Begin assisted forward elevation (FE).
- Perform pulley exercises and teach for home program.
- Perform isometric exercises for IR, ER, extension, and abduction.

Phase III (7 weeks – 2 months)

- Begin supine active FE.
- Progressively increase patient's position from supine to erect during FE exercises.
- Use therabands of progressive strengths for IR, ER, flexion, abduction, and extension. Goal is progressive strengthening of deltoid and rotator cuff.

Begin flexibility and stretching exercises to progressively increase