



James S. Kercher, M.D.  
Sports Medicine, Shoulder &  
Knee Surgery, Cartilage Restoration

## **Unstable Proximal Humerus Fracture**

**Goal:** Maintain ROM without displacing the proximal humerus fracture.

### **Phase I (0 to 21 days)**

- Begin elbow, wrist and hand active ROM.
- A sling and swath, or Velpeau with an axillary pad should be used when not doing physical therapy.
- After 14 days, begin pendulum exercises (clockwise and counterclockwise).
- Establish a home exercise program so patient is performing exercises 3-5 times per day for 10 minutes each session.

### **Phase II (3 to 6 weeks)**

- Begin supine ER with a cane. Fifteen to 20 degrees of abduction are permitted if the patient is more comfortable.
- Begin active assisted forward elevation (FE).
- Perform pulley exercises and teach for home program.
- Perform isometric exercises for IR, ER, extension, and abduction.

### **Phase III (7 weeks to 2 months)**

- Begin supine active FE. Progressively increase patient's position from supine to erect during FE exercises.
- Use Therabands of progressive strengths for IR, ER, anterior, middle and posterior deltoid.

Begin flexibility and stretching exercises to progressively increase ROM in all positions (i.e. towel behind back, finger walking up the wall, etc.).